

Homoclinic, Subharmonic and Superharmonic Bifurcations for a Pendulum with Periodically Varying Length

A. O. Belyakov

*Institute of Mechanics, Lomonosov Moscow State University,
Michurinsky pr. 1, Moscow, Russia
ORCOS, Institute of Mathematical Methods in Economics,
Vienna University of Technology
Argentinierstrasse 8/105-4, A-1040 Vienna, Austria*

A. P. Seyranian

*Institute of Mechanics, Lomonosov Moscow State University,
Michurinsky pr. 1, Moscow, Russia*

Abstract

Dynamic behavior of a weightless rod with a point mass sliding along the rod axis according to periodic law is studied. This is the simplest model of child's swing. Melnikov's analysis is carried out to find bifurcations of homoclinic, subharmonic oscillatory, and subharmonic rotational orbits. For the analysis of superharmonic rotational orbits the averaging method is used. The analytical results are compared with numerical simulation results.

1 Introduction

Oscillations of a pendulum with periodically varying length (PPVL) is one of the classical problems in mechanics, see [1–10]. We represent PPVL as a weightless rod with a point mass sliding along the rod axis according to periodic law. This is also a simple model of child's swing. In works [1,6–9] small oscillations of PPVL were studied. In [5,8,9] the instability domains of

Email addresses: belyakov@imec.msu.ru, anton.belyakov@tuwien.ac.at (A. O. Belyakov), seyran@imec.msu.ru (A. P. Seyranian).

the vertical position were found both analytically and numerically. Regular rotations and chaotic regimes were also investigated in [6,8,9].

In the literature oscillatory and rotational approximate solutions for PPVL were obtained with quasi-linear approach, where nonlinearity was assumed to be small as well as the excitation amplitude. The only exception is [10], where stable uniform rotations were found in the case of zero damping and special excitations.

In the present paper we analyze boundaries in the parameter space for oscillatory, rotational, and more complex (rotational-oscillatory, chaotic) regimes of PPVL assuming arbitrary (not small) nonlinearity. In Section 2 main equations of motion of a pendulum with variable length are derived and given in non-dimensional form. In Section 3 we use Melnikov's analysis [11] to find bifurcations of homoclinic, oscillatory and subharmonic rotational orbits of PPVL. Melnikov's functions are obtained and compared with the results of numerical simulation. Similar analysis was done for a pendulum with vibrating pivot in [14]. In Section 4 we find bifurcations of superharmonic rotational orbits with the use of the averaging method and compare them with the numerical simulation results. The methodological novelty in Section 4 is that the unperturbed system preserves the angular momentum (rather than Hamiltonian as in the Melnikov's analysis) and allows for non-small excitation amplitudes.

2 Main relations

Equation for motion of the PPVL is derived with the use of angular momentum alteration theorem and taking into account linear damping forces, see [5,6,8]

$$\frac{d}{dt} \left(ml^2 \frac{d\theta}{dt} \right) + \gamma l^2 \frac{d\theta}{dt} + mgl \sin(\theta) = 0, \quad (1)$$

where m is the mass, l is the length, θ is the angle of the pendulum deviation from the vertical position, γ is the damping coefficient, and g is the acceleration due to gravity.

It is assumed that the length of the pendulum changes according to the periodic law

$$l = l_0 + a\varphi(\Omega t), \quad (2)$$

where l_0 is the mean pendulum length, a and Ω are the amplitude and frequency of the excitation, $\varphi(\tau)$ is the smooth zero mean periodic function with

period 2π .

We introduce new time $\tau = \Omega t$ and three dimensionless parameters

$$\varepsilon = \frac{a}{l_0}, \quad \omega = \frac{\Omega_0}{\Omega}, \quad \beta = \frac{\gamma}{m\Omega_0}, \quad (3)$$

where $\Omega_0 = \sqrt{\frac{g}{l_0}}$ is the eigenfrequency of the pendulum with constant length $l = l_0$ and zero damping. In this notations equation (1) takes the form

$$\left((1 + \varepsilon\varphi(\tau))^2 \dot{\theta} \right)' + \beta\omega (1 + \varepsilon\varphi(\tau))^2 \dot{\theta} + (1 + \varepsilon\varphi(\tau)) \omega^2 \sin(\theta) = 0, \quad (4)$$

where the upper dot denotes differentiation with respect to new time τ .

3 Melnikov's method: Perturbation of a Hamiltonian System

Assuming $1 + \varepsilon\varphi(\tau) > 0$, equation (4) can be written in the following form

$$\ddot{\theta} + \left(\frac{2\varepsilon\dot{\varphi}(\tau)}{1 + \varepsilon\varphi(\tau)} + \beta\omega \right) \dot{\theta} + \frac{\omega^2 \sin(\theta)}{1 + \varepsilon\varphi(\tau)} = 0. \quad (5)$$

Coefficients of nonlinear equation (5) explicitly depend on periodic function $\varphi(\tau)$ and three independent dimensionless parameters: the relative excitation amplitude ε , the damping β , and the inverse relative frequency of excitation ω .

Let us assume that parameters of excitation amplitude ε and damping β are small of the same order, $\varepsilon \sim \beta \ll 1$. Thus, we can say that dynamics of PPVL is described by the perturbed Hamiltonian system

$$\dot{\theta} = \frac{\partial H}{\partial v}, \quad (6)$$

$$\dot{v} = -\frac{\partial H}{\partial \theta} + g_1(\theta, v, \tau) + o(\varepsilon), \quad (7)$$

where the perturbation function

$$g_1(\theta, v, \tau) = (2\varepsilon \sin(\tau) - \beta\omega) v + \varepsilon\omega^2 \cos(\tau) \sin(\theta) \quad (8)$$

is of the first order of smallness, i.e. $g_1(\theta, v, \tau) = O(\varepsilon)$. The following function

$$H = \frac{v^2}{2} - \omega^2 \cos(\theta) \quad (9)$$

is the Hamiltonian of the unperturbed system

$$\dot{\theta} = v, \quad (10)$$

$$\dot{v} = -\omega^2 \sin(\theta), \quad (11)$$

which is system (6)–(8) with $\varepsilon = 0$ and $\beta = 0$. The unperturbed system (10)–(11) describes the oscillations of the pendulum with constant length and zero damping, so that the integral $H = \text{const}$.

Unperturbed system (6)–(8) has an oscillatory solution if $H < \omega^2$ and a rotational solution if $H > \omega^2$. If $H = \omega^2$ the solution is a separatrix dividing oscillatory and rotational domains in phase space.

3.1 Homoclinic bifurcations

In order to apply Melnikov's criterion [11,12] to a homoclinic orbit we find the separatrix of the unperturbed system (10)–(11) that goes through the saddle-node point $\theta = \pi$, $v = 0$. For this point we have $H = \omega^2$ and with the use of (9) obtain $v^2 = 2\omega^2(1 + \cos(\theta)) = 4\omega^2 \cos(\theta/2)$. So, the separatrix has the following form

$$v = \dot{\theta} = \pm 2\omega \cos\left(\frac{\theta}{2}\right). \quad (12)$$

This equation allows for separation of variables and the following integration

$$\ln\left(\frac{1 + \sin(\theta/2)}{\cos(\theta/2)}\right) = \pm\omega(\tau - \tau_0), \quad (13)$$

where τ_0 is a constant of integration. Potentiation of (13) with some transformations yields

$$\cos\left(\frac{\theta}{2}\right) = \frac{1}{\cosh(\omega(\tau - \tau_0))}, \quad \sin\left(\frac{\theta}{2}\right) = \tanh(\omega(\tau - \tau_0)). \quad (14)$$

Melnikov's distance between stable and unstable perturbed separatrices is given by the following integral

$$\begin{aligned}
M^\pm &= \int_{-\infty}^{\infty} \frac{\partial H}{\partial v} g_1(\theta, v, \tau) d\tau \\
&= \int_{-\infty}^{\infty} \left((2\varepsilon \sin(\tau) - \beta\omega) v^2(\tau) + \varepsilon\omega^2 v(\tau) \cos(\tau) \sin(\theta(\tau)) \right) d\tau
\end{aligned}$$

which with the use of (12) and (14) takes the form

$$\begin{aligned}
M^\pm &= 8\varepsilon\omega^2 \int_{-\infty}^{\infty} \frac{\sin(\tau) d\tau}{\cosh^2(\omega(\tau - \tau_0))} - 4\beta\omega^3 \int_{-\infty}^{\infty} \frac{d\tau}{\cosh^2(\omega(\tau - \tau_0))} \\
&\quad + 4\varepsilon\omega^3 \int_{-\infty}^{\infty} \frac{\cos(\tau) \sinh(\omega(\tau - \tau_0))}{\cosh^3(\omega(\tau - \tau_0))} d\tau,
\end{aligned}$$

and has the following expression (see Appendix A)

$$M^\pm = \frac{6\pi\varepsilon \sin(\tau_0)}{\sinh\left(\frac{\pi}{2\omega}\right)} - 8\beta\omega^2. \quad (15)$$

This is a sign-changing function if the following inequality is satisfied

$$\frac{\varepsilon}{\beta} > \frac{4\omega^2}{3\pi} \sinh\left(\frac{\pi}{2\omega}\right). \quad (16)$$

According to Melnikov's criterion [11] when condition (16) is satisfied then homoclinic structure and complex dynamics appear near the separatrix. Condition (16) means that complex dynamics appears only when the amplitude of excitation is sufficiently large with respect to the damping coefficient. The domain of possible complex dynamics defined by inequality (16) is depicted in Fig 1 and compared with numerical simulations in Fig 2 for $\beta = 0.05$. The minimum of the right-hand side of (16) is reached at $\omega \approx 0.82$ and can be evaluated as 0.948. Note that condition (16) is similar to that of oscillator with quadratic nonlinearity and external periodic excitation [13]. Similar inequality for the pendulum with vertically vibrating pivot was obtained in [14], where the right-hand side function is three times greater than that in (16).

3.2 Subharmonic bifurcations of oscillatory orbits

In order to apply Melnikov's criterion to oscillatory orbits we find oscillatory solutions of the unperturbed system (10)-(11). For this reason we introduce the amplitude of oscillations A so that from the first integral for $v = 0$ we have $H =$

$-\omega^2 \cos(A)$ and with the use of (9) we obtain $v^2 = 2\omega^2 (\cos(\theta) - \cos(A)) = 4\omega^2 (\sin^2(A/2) - \sin^2(\theta/2))$. Thus, we can write the following equation

$$v = \dot{\theta} = 2\omega \operatorname{sign}(\dot{\theta}) \sqrt{\sin^2\left(\frac{A}{2}\right) - \sin^2\left(\frac{\theta}{2}\right)}, \quad (17)$$

which allows for separation of variables. In order to integrate (17) one usually introduces a monotonically increasing phase ψ such that

$$\sin\left(\frac{\theta}{2}\right) = k \sin(\psi),$$

where new variable $k = \sin\left(\frac{A}{2}\right) = \sqrt{\frac{H+\omega^2}{2\omega^2}}$ is called modulus in elliptic integrals. Skipping intermediate transformations (see [15]) we have $\psi = \operatorname{am}(\omega(\tau - \tau_0), k)$ and consequently

$$v = 2\omega k \operatorname{cn}(\omega(\tau - \tau_0), k) \quad (18)$$

and

$$\cos\left(\frac{\theta}{2}\right) = \operatorname{dn}(\omega(\tau - \tau_0), k), \quad \sin\left(\frac{\theta}{2}\right) = k \operatorname{sn}(\omega(\tau - \tau_0), k), \quad (19)$$

where am , dn , cn , and sn are the Jacobi functions. The k value follows from the resonance condition stating that period of oscillation $\frac{4K(k)}{\omega}$ and period of excitation 2π should be in rational relation

$$\frac{4K(k)}{\omega} p = 2\pi q, \quad (20)$$

where p and q are relatively prime natural numbers and $K(k)$ is the complete elliptic integral of the first kind. Thus, for oscillation motion in resonance $p:q$ we have the following subharmonic Melnikov's distance with the use of expressions (18), (19)

$$\begin{aligned} M^{q/p} &= \int_0^{2\pi q} \frac{\partial H}{\partial v} g_1(\theta, v, \tau) d\tau \\ &= \int_0^{2\pi q} \left((2\varepsilon \sin(\tau) - \beta\omega)v^2(\tau) + \varepsilon\omega^2 v(\tau) \cos(\tau) \sin(\theta(\tau)) \right) d\tau \end{aligned}$$




Fig. 1. Subharmonic bifurcation functions (ε/β) which is the boundary in (23) for $q = 2, q = 4, q = 6$, and $p = 1$ converge to the homoclinic bifurcation function that approximates the domain of complex dynamics in (16).

$$\begin{aligned}
&= 4\omega^2 k^2 \int_0^{2\pi q} (2\varepsilon \sin(\tau) - \beta\omega) \operatorname{cn}^2(\omega(\tau - \tau_0), k) d\tau \\
&\quad + 4\varepsilon\omega^3 k^2 \int_0^{2\pi q} \cos(\tau) \operatorname{sn}(\omega(\tau - \tau_0), k) \operatorname{dn}(\omega(\tau - \tau_0), k) \operatorname{cn}(\omega(\tau - \tau_0), k) d\tau.
\end{aligned} \tag{21}$$

Taking the integrals in Appendix B for $p = 1$ and even $q = 2, 4, 6, \dots$ we have the following Melnikov's distance

$$M^{1/q} = 4\omega^4 \left(\frac{3\varepsilon\pi}{\omega^2} \frac{\sin \tau_0}{\sinh(K(k')/\omega)} - 4\beta \left(E(k) - (k'^2)K(k) \right) \right), \tag{22}$$

where $k'^2 = 1 - k^2$. This is a sign-changing function if the following inequality is satisfied

$$\frac{\varepsilon}{\beta} > \frac{4\omega^2}{3\pi} \left(E(k) - (k'^2)K(k) \right) \sinh\left(\frac{K(k')}{\omega}\right). \tag{23}$$

Condition (23) means that corresponding $1:q$ resonant oscillations appear only when the amplitude of excitation is sufficiently large with respect to the damping coefficient. The domain of possible oscillations defined by inequality (23) is depicted in Fig 1 and compared with numerical simulations in Fig 2 for $\beta = 0.05$. Note that condition (23) is similar to that of oscillator with quadratic




Fig. 2. Periods of oscillations divided by period of excitation ($p : q$) are shown with different colors on the parameter plane (ω, ε) at $\beta = 0.05$. The correspondence between the colors and relative periods is shown by the color bar on the right, where stationary and rotational regimes are shown with white color, while chaotic regimes – with black. Subharmonic bifurcation functions for $q = 2$ (solid line), $q = 4$ (dot-and-dash line), $q = 6$ (dotted line), and $p = 1$ converge to the homoclinic bifurcation function (bold solid line) and approximate the domains where corresponding resonant oscillations can occur, shown with blue ($q = 2$), purple ($q = 4$), and red ($q = 6$) colors.

nonlinearity and external periodic excitation [13]. Similar inequality for the pendulum with vertically vibrating pivot was obtained in [14], where the right-hand side function is three times greater than that in (23).

3.3 Subharmonic bifurcations of rotational orbits

In order to apply Melnikov's criterion to a rotational orbit we find the solution of the unperturbed system (10)–(11) for $H > \omega^2$. Thus, with the use of (9) we obtain $\dot{\theta} = \pm\sqrt{2}\sqrt{H + \omega^2 \cos(\theta)} = \pm 2\omega k \sqrt{1 - \sin^2(\theta/2)/k^2}$, where “ $\pm$ ” represents counter- and clockwise rotations. Since the model is symmetric with respect to the vertical axis, we will consider only the counterclockwise rotation (“+” instead of “ $\pm$ ”). This solution has the form $\theta = 2 \operatorname{am}(\omega k (\tau - \tau_0), 1/k)$, so we have

$$v = \dot{\theta} = 2\omega k \operatorname{dn}\left(\omega k (\tau - \tau_0), \frac{1}{k}\right), \quad (24)$$

$$\cos\left(\frac{\theta}{2}\right) = \operatorname{cn}\left(\omega k(\tau - \tau_0), \frac{1}{k}\right), \quad \sin\left(\frac{\theta}{2}\right) = \operatorname{sn}\left(\omega k(\tau - \tau_0), \frac{1}{k}\right). \quad (25)$$

The k value follows from the resonance condition stating that period of rotation $\frac{2K(1/k)}{\omega k}$ and period of excitation 2π should be in rational relation

$$\frac{2K\left(\frac{1}{k}\right)}{\omega k} r = 2\pi q, \quad (26)$$

where r and q are relatively prime natural numbers. Thus, for rotational motion in resonance $r:q$ we have the following subharmonic Melnikov's distance

$$\begin{aligned} M^{q/r} &= \int_0^{2\pi q} \frac{\partial H}{\partial v} g_1(\theta, v, \tau) d\tau \\ &= \int_0^{2\pi q} \left((2\varepsilon \sin(\tau) - \beta\omega) v^2(\tau) + \varepsilon\omega^2 v(\tau) \cos(\tau) \sin(\theta(\tau)) \right) d\tau \\ &= 8\varepsilon\omega^2 k^2 \int_0^{2\pi q} (2\varepsilon \sin(\tau) - \beta\omega) \operatorname{dn}^2\left(\omega k(\tau - \tau_0), \frac{1}{k}\right) d\tau \\ &\quad + 4\varepsilon\omega^3 k \int_0^{2\pi q} \cos(\tau) \operatorname{sn}\left(\omega k(\tau - \tau_0), \frac{1}{k}\right) \operatorname{dn}\left(\omega k(\tau - \tau_0), \frac{1}{k}\right) \operatorname{cn}\left(\omega k(\tau - \tau_0), \frac{1}{k}\right) d\tau. \end{aligned}$$

Taking the integrals in Appendix C for $r = 1$ and $q = 1, 2, 3, \dots$ we have the following Melnikov's distance

$$M^{q/1} = -2\omega^2 \left(\frac{3\varepsilon\pi}{\omega^2} \frac{\sin \tau_0}{\sinh\left(\frac{K'}{\omega k}\right)} - 4\beta k E\left(\frac{1}{k}\right) \right), \quad (27)$$

where $K' = K(\sqrt{1 - 1/k^2})$. This is a sign-changing function if the following inequality is satisfied

$$\frac{\varepsilon}{\beta} > \frac{4\omega^2 k}{3\pi} E\left(\frac{1}{k}\right) \sinh\left(\frac{K'}{\omega k}\right). \quad (28)$$

Condition (28) means that corresponding $1:q$ resonant rotations appear only when the amplitude of excitation is sufficiently large with respect to the damping coefficient. The domain of possible rotations defined by inequality (28) is depicted in Fig. 3 and compared with numerical simulations in Fig. 4 for




Fig. 3. Subharmonic bifurcation functions (ε/β) which is the boundary in (28) for $q = 1$, $q = 2$, $q = 3$, and $r = 1$ converge to the homoclinic bifurcation function that approximates the domain of complex dynamics in (16).

$\beta = 0.05$. Although numerically found rotations 1:2 are not monotone. These are rotation-oscillation regimes. Note that condition (28) is similar to that of oscillator with quadratic nonlinearity and external periodic excitation [13]. Similar inequality for the pendulum with vertically vibrating pivot was obtained in [14], where the right-hand side function is three times greater than that in (28).

4 Averaging method: Superharmonic bifurcations of rotational orbits

We study resonant rotations $r:q = 1:1, 2:1, 3:1, \dots$ under assumption that ω is small (e.g., because of the small gravitation g or high excitation frequency Ω) of the same order as damping, $\beta \sim \omega \ll 1$, while the excitation amplitude ε is not small. Then, unperturbed system (equation (4) with $\beta = \omega = 0$) has the specific angular momentum or sector velocity $s = (1 + \varepsilon\varphi(\tau))^2 \dot{\theta}$ as its first integral. The unperturbed system is not Hamiltonian, so we cannot apply Melnikov's analysis to obtain domains of existence of corresponding rotations. For this purpose we use the method of averaging, [2,17]. Equation (4) can be written in the form of the system




Fig. 4. Periods of regular rotations divided by period of excitation (this ratio is $r : q$ for oscillations) are shown with different colors on the parameter plane (ω, ε) at $\beta = 0.05$. The correspondence between the colors and relative periods q is shown by the color bar on the right, where stationary and chaotic regimes are shown with white color. Subharmonic bifurcation functions for $q = 1$, $q = 2$, $q = 3$, and $r = 1$ converge to the homoclinic bifurcation function and approximate the domains where corresponding resonant rotations can occur.

$$\dot{\theta} = \frac{s}{(1 + \varepsilon\varphi(\tau))^2}, \quad (29)$$

$$\dot{s} = f(\theta, s, \tau), \quad (30)$$

where the perturbation function

$$f(\theta, s, \tau) = -\beta\omega s - (1 + \varepsilon\varphi(\tau))\omega^2 \sin(\theta) \quad (31)$$

is of the second order of smallness, meaning $f(\theta, p, \tau) = O(\omega^2)$. The unperturbed system has the solution

$$\theta = s \int_0^\tau \frac{d\eta}{(1 + \varepsilon\varphi(\eta))^2} + \vartheta = s \Phi(\tau, \varepsilon) + \vartheta, \quad (32)$$

where $\Phi(\tau, \varepsilon) = \int_0^\tau \frac{d\eta}{(1 + \varepsilon\varphi(\eta))^2}$ and ϑ is the constant phase shift. In the first approximation we can derive the averaged equation of (30) substituting in it

expression (32) and taking the average of its right-hand side, see [17]

$$\dot{\bar{s}} = \frac{1}{2\pi q r} \int_0^{2\pi q r} f(\bar{s} \Phi(\tau, \varepsilon) + \bar{\vartheta}, \bar{s}, \tau) d\tau \quad (33)$$

where the integral is taken as if corresponding *averaged variables* $\bar{s}$ and $\bar{\vartheta}$ are constant. The steady state value $\bar{s}$ follows from the resonance condition written with the use of solution (32) and stating that period of rotation and period of excitation should be in rational relation

$$2\pi r = \bar{s} \int_0^{2\pi q} \frac{d\tau}{(1 + \varepsilon \varphi(\tau))^2} = \bar{s} q \Phi(2\pi, \varepsilon), \quad (34)$$

where r and q are relatively prime natural numbers. Steady state ($\dot{\bar{s}} = 0$) value of $\bar{\vartheta}$ we find from (33).¹ Thus, with the use of (34), for rotational motion in resonance $r:q$ we have

$$\begin{aligned} -\frac{(2\pi r)^2 \beta}{\omega \Phi(2\pi, \varepsilon)} &= \int_0^{2\pi r q} (1 + \varepsilon \varphi(\tau)) \sin\left(2\pi \frac{r}{q} \frac{\Phi(\tau, \varepsilon)}{\Phi(2\pi, \varepsilon)} + \bar{\vartheta}\right) d\tau \\ &= A(\varepsilon, r, q) \cos(\bar{\vartheta}) + B(\varepsilon, r, q) \sin(\bar{\vartheta}) \end{aligned} \quad (35)$$

where $A(\varepsilon, r, q)$ and $B(\varepsilon, r, q)$ denote the following integrals

$$A(\varepsilon, r, q) = \int_0^{2\pi r q} (1 + \varepsilon \varphi(\tau)) \sin\left(2\pi \frac{r}{q} \frac{\Phi(\tau, \varepsilon)}{\Phi(2\pi, \varepsilon)}\right) d\tau, \quad (36)$$

$$B(\varepsilon, r, q) = \int_0^{2\pi r q} (1 + \varepsilon \varphi(\tau)) \cos\left(2\pi \frac{r}{q} \frac{\Phi(\tau, \varepsilon)}{\Phi(2\pi, \varepsilon)}\right) d\tau. \quad (37)$$

¹ One can also write the differential equation for slow phase shift as $\dot{\vartheta} = \Phi(\tau, \varepsilon) f(s \Phi(\tau, \varepsilon) + \vartheta, s, \tau)$, but it has unbounded right hand side because of the growing multiplier $\Phi(\tau, \varepsilon) \sim \tau$. That is why this equation does not fit for standard Bogolubov's method of averaging, see [2]. Thus, we can obtain here only the first order approximation with the use of averaged equation (33). In order to obtain higher order approximations the general averaging scheme by Volosov can be applied to system (29)-(30), see [17].




Fig. 5. Superharmonic bifurcation functions (ω/β) which is the boundary in (38) for $r = 1$, $r = 2$, $r = 3$, and $q = 1$.

Thus, the domain of existence of corresponding regular rotations is approximated by the following condition that equation (35) has a solution

$$\frac{\omega}{\beta} \geq \frac{(2\pi r)^2}{\Phi(2\pi, \varepsilon) \sqrt{A^2(\varepsilon, r, q) + B^2(\varepsilon, r, q)}}, \quad (38)$$

which right hand side can be calculated numerically for any particular ε , r , and q , see Fig. 5, where $\varphi(\tau) = \cos(\tau)$.

In order to find the boundaries for rotational regimes $r = 1, 2, 3$ the first approximation is enough because excitation amplitude ε is not small in contrast to the quasilinear approach in [9], where both ε and ω are assumed to be small and higher order approximations of the averaging method are needed to obtain similar boundaries.

5 Conclusion

For the pendulum with variable length we derived analytical formulas for the boundaries of bifurcations in the space of three parameters: relative frequency of excitation, amplitude of excitation, and damping. The boundaries for homoclinic bifurcation separating the domain of only stationary and oscillatory




Fig. 6. Periods of regular rotations divided by period of excitation (this ratio is $r : q$ for oscillations) are shown with different colors on the parameter plane (ω, ε) at $\beta = 0.05$. The correspondence between the colors and relative periods r is shown by the color bar on the right, where stationary, oscillatory, and chaotic regimes are shown with white color. Superharmonic bifurcation functions for $r = 1$, $r = 2$, $r = 3$, and $r = 1$ bound from below corresponding regimes obtained numerically.

regimes from the domain of more complex dynamics, subharmonic oscillations and subharmonic rotations are obtained using Melnikov's method under assumption of small damping and excitation amplitude. For the analysis of superharmonic bifurcations of rotational orbits the method of averaging is used assuming smallness of relative excitation frequency rather than that of excitation amplitude. Both methods allow to obtain in the first approximation the basic rotational orbit with angular velocity equal to the excitation frequency, 1:1. Small excitation frequency (or small gravity) allows to introduce the unperturbed system with the conservation of angular momentum, so that faster rotations are found in the first approximation by the method of averaging. In Figs. 2, 4, and 6 it is shown that the boundaries for complex dynamics, subharmonic oscillations, and superharmonic rotations are in good agreement with the results of numerical simulation.

Appendices

A Melnikov function for homoclinic orbit

$M^\pm = 8\varepsilon\omega^2 I_1 - 4\beta\omega^3 I_2 + 4\varepsilon\omega^3 I_3$, where the first integral

$$\begin{aligned}
I_1 &= \int_{-\infty}^{\infty} \frac{\sin(\tau) d\tau}{\cosh^2(\omega(\tau - \tau_0))} \\
&= \frac{1}{\omega} \int_{-\infty}^{\infty} \frac{\sin(\tau_0 + \eta/\omega) d\eta}{\cosh^2(\eta)} \\
&= \frac{\sin(\tau_0)}{\omega} \int_{-\infty}^{\infty} \frac{\cos(\eta/\omega) d\eta}{\cosh^2(\eta)} + \frac{\cos(\tau_0)}{\omega} \int_{-\infty}^{\infty} \frac{\sin(\eta/\omega) d\eta}{\cosh^2(\eta)}. \tag{A.1}
\end{aligned}$$

The integral $\int_{-\infty}^{\infty} \frac{\sin(\eta/\omega) d\eta}{\cosh^2(\eta)}$ is zero because its integrand is an odd function, while the other integral has even integrand and can be calculated as follows $\int_{-\infty}^{\infty} \frac{\cos(t/\omega)}{\cosh^2 t} dt = \frac{\pi}{\omega \sinh(\pi/2\omega)}$, hence, the first term has the expression

$$I_1 = \frac{\pi \sin(\tau_0)}{\omega^2 \sinh(\pi/2\omega)}. \tag{A.2}$$

The second integral can be calculated as follows

$$I_2 = \int_{-\infty}^{\infty} \frac{d\tau}{\cosh^2(\omega(\tau - \tau_0))} = \frac{1}{\omega} \int_{-\infty}^{\infty} \frac{ds}{\cosh^2(s)} = \frac{2}{\omega}, \tag{A.3}$$

while the third integral can be converted to the first one via integration by parts using the relation $\frac{\sinh(s) ds}{\cosh^3(s)} = -\frac{1}{2} d\frac{1}{\cosh^2(s)}$

$$\begin{aligned}
I_3 &= \int_{-\infty}^{\infty} \frac{\cos(\tau) \sinh(\omega(\tau - \tau_0))}{\cosh^3(\omega(\tau - \tau_0))} d\tau, \\
&= \frac{1}{\omega} \int_{-\infty}^{\infty} \frac{\cos(\tau_0 + \eta/\omega) \sinh(\eta)}{\cosh^3(\eta)} d\eta \\
&= -\frac{1}{2\omega^2} \left. \frac{\cos(\tau_0 + \eta/\omega)}{\cosh^2(\eta)} \right|_{-\infty}^{\infty} - \frac{1}{2\omega^2} \int_{-\infty}^{\infty} \frac{\sin(\tau_0 + \eta/\omega)}{\cosh^2(\eta)} d\eta = -\frac{I_1}{2\omega}. \tag{A.4}
\end{aligned}$$

Thus we have Melnikov function $M^\pm = 8\varepsilon\omega^2 I_1 - 4\beta\omega^3 I_2 + 4\varepsilon\omega^3 I_3 = \frac{6\pi\varepsilon \sin(\tau_0)}{\sinh(\pi/2\omega)} - 8\beta\omega^2$.

B Melnikov function for subharmonic oscillations

$M^{q/p} = 8\varepsilon\omega^2 k^2 I_1 - 4\beta\omega^3 k^2 I_2 + 4\varepsilon\omega^3 k^2 I_3$, where

$$\begin{aligned} I_1 &= \int_0^{2\pi q} \sin(\tau) \operatorname{cn}^2(\omega(\tau - \tau_0), k) \, d\tau \\ &= -\cos(\tau) \operatorname{cn}^2(\omega(\tau - \tau_0), k) \Big|_0^{2\pi q} - 2\omega I_3, \\ &= -2\omega I_3, \end{aligned} \tag{B.1}$$

$$I_2 = \int_0^{2\pi q} \operatorname{cn}^2(\omega(\tau - \tau_0), k) \, d\tau, \tag{B.2}$$

$$I_3 = \int_0^{2\pi q} \cos(\tau) \operatorname{cn}(\omega(\tau - \tau_0), k) \operatorname{sn}(\omega(\tau - \tau_0), k) \operatorname{dn}(\omega(\tau - \tau_0), k) \, d\tau \tag{B.3}$$

where we use the formula $\frac{d \operatorname{cn}(u)}{du} = -\operatorname{sn}(u) \operatorname{dn}(u)$. Thus, we have $M^{q/p} = -12\varepsilon\omega^3 k^2 I_3 - 4\beta\omega^3 k^2 I_2$.

$$\begin{aligned} I_1 &= \int_0^{2\pi q} \sin(\tau) \operatorname{cn}^2(\omega(\tau - \tau_0), k) \, d\tau \\ &= \frac{1}{\omega} \int_0^{2\pi q\omega} \sin(\tau_0 + s/\omega) \operatorname{cn}^2(s, k) \, ds \\ &= \frac{\sin(\tau_0)}{\omega} \int_0^{2\pi q\omega} \cos(s/\omega) \operatorname{cn}^2(s, k) \, ds \end{aligned} \tag{B.4}$$

The integral I_3 vanishes except for $p = 1$ and even q . In this case we have from (312.02) in [16]

$$I_2 = \frac{4\omega}{k^2} \left(E(k) - (k'^2) K(k) \right),$$

$$I_3 = -\frac{\pi}{k^2\omega} \frac{\sin \tau_0}{\sinh(K(k'))},$$

where $k'^2 = 1 - k^2$. So we have $M^{q/1} = 4\omega^4 \left(\frac{3\varepsilon\pi}{\omega^2} \frac{\sin \tau_0}{\sinh(K(k'))} - 4\beta (E(k) - (k'^2) K(k)) \right)$.

C Melnikov function for subharmonic rotations

$$M^{q/r} = 8\varepsilon\omega^2 k^2 I_1 - 4\beta\omega^3 k^2 I_2 + 4\varepsilon\omega^3 k I_3,$$

$$\begin{aligned} I_1 &= \int_0^{2\pi q} \sin(\tau) \operatorname{dn}^2\left(\omega k (\tau - \tau_0), \frac{1}{k}\right) d\tau \\ &= -\cos(\tau) \operatorname{dn}^2\left(\omega k (\tau - \tau_0), \frac{1}{k}\right) \Big|_0^{2\pi q} - \frac{2\omega}{k} I_3 = -\frac{2\omega}{k} I_3, \end{aligned} \quad (\text{C.1})$$

$$I_2 = \int_0^{2\pi q} \operatorname{dn}^2\left(\omega k (\tau - \tau_0), \frac{1}{k}\right) d\tau, \quad (\text{C.2})$$

$$I_3 = \int_0^{2\pi q} \cos(\tau) \operatorname{sn}\left(\omega k (\tau - \tau_0), \frac{1}{k}\right) \operatorname{dn}\left(\omega k (\tau - \tau_0), \frac{1}{k}\right) \operatorname{cn}\left(\omega k (\tau - \tau_0), \frac{1}{k}\right) d\tau, \quad (\text{C.3})$$

where we use the formula $\frac{d \operatorname{dn}(u, 1/k)}{du} = -\operatorname{sn}(u, 1/k) \operatorname{cn}(u, 1/k)/k^2$. Thus, we have $M^{q/r} = -12\varepsilon\omega^3 k^2 I_3 - 4\beta\omega^3 k^2 I_2$.

The integral I_3 vanishes except for $r = 1$. In this case we have

$$I_2 = \frac{2}{\omega k} E\left(\frac{1}{k}\right),$$

$$I_3 = -\frac{\pi}{k^2 \omega} \frac{\sin \tau_0}{\sinh(K')},$$

where $K' = K\left(\sqrt{1 - 1/k^2}\right)$. So we have $M^{q/1} = -2\omega^2 \left(\frac{3\varepsilon\pi}{\omega^2} \frac{\sin \tau_0}{\sinh\left(\frac{K'}{\omega k}\right)} - 4\beta E\left(\frac{1}{k}\right) \right)$.

References

- [1] H. Kauderer, *Nichtlineare Mechanik*, Springer, Berlin, 1958.
- [2] N.N. Bogoliubov, Y.A. Mitropolsky, *Asymptotic Methods in the Theory of Non-Linear Oscillations*, Gordon and Breach, New York, 1961.
- [3] Ya.G. Panovko, I.I. Gubanova, *Stability and Oscillations of Elastic Systems. Modern Concepts, Paradoxes and Mistakes*, Nauka, Moscow, 1987.
- [4] K. Magnus, *Schwingungen. Eine Einfuhrung in die theoretische Behandlung von Schwingungsproblemen*, J.Teubner, Stuttgart, 1976.
- [5] A.P. Seyranian, The swing: parametric resonance, *Journal of Applied Mathematics and Mechanics* 68 (2004) 757–764.

- [6] A.P. Seyranian, A.O. Belyakov, Swing dynamics, *Doklady Physics* 53 (2004) 388–394.
- [7] A.A. Zevin, L.A. Filonenko, Qualitative study of oscillations of a pendulum with periodically varying length and a mathematical model of swing, *Journal of Applied Mathematics and Mechanics*, 71 (2007) 989–1003.
- [8] A.O. Belyakov, A.P. Seyranian, A. Luongo, Dynamics of the pendulum with periodically varying length, *Physica D* 238 (2009) 1589–1597.
- [9] A.O. Belyakov, A.P. Seyranian, On Nonlinear Dynamics of the Pendulum with Periodically Varying Length, *Mechanics of Machines, BulKToMM*, 18, No 3 (2010) 21–28, (ISSN 0861-9727), arXiv:0910.3802v2 [math-ph].
- [10] A.P. Markeev, Uniform rotations of a variable-length pendulum, *Doklady Physics*, 56 (2011) 240–243.
- [11] V.K. Melnikov, On stability of a center under periodic in time perturbations, *Trudy Moskovskogo Matematicheskogo Obshchestva* 12 (1963) 3–52.
- [12] J. Guckenheimer, P.J. Holmes, *Nonlinear Oscillations, Dynamical Systems and Bifurcations of Vector Fields*, Springer, Berlin, 1983.
- [13] S.P. Kuznetsov, *Dynamical Chaos*, Physmatlit, Moscow, 2006.
- [14] B.P. Koch, R.W. Leven, Subharmonic and homoclinic bifurcations in a parametrically forced pendulum, *Physica D: Nonlinear Phenomena*, 16 (1985) 1–13.
- [15] Ch.J. de La Vallée Poussin, *Leçon de Mécanique Analytique*. Vol. 1., Leuven, 1932.
- [16] P.F. Byrd, M.D. Friedman, *Handbook of elliptic integrals for engineers and scientists*, second ed., Springer, 1971.
- [17] V.M. Volosov, B.I. Morgunov, *Averaging Method in the Theory of Nonlinear Oscillatory Systems*, MSU, Moscow, 1971.